

# Advancing health equity by addressing social determinants of health: Using health data to improve educational outcomes

Mary Jo Rattermann, Azure Angelov, Tommy Reddicks, Jess Monk

Published: March 17, 2021 • <https://doi.org/10.1371/journal.pone.0247909>

## Abstract

Data from two social determinants of health—access to health care and access to a quality education—are combined to examine the impact of health on student achievement. Data from a high poverty, high performing K-8 school revealed a significant negative correlation between the number of visits to a school-based nurse and standardized academic assessments. Fixed effect regression confirmed the effect of total number of visits to the school-based nurse on performance on standardized assessments, and also revealed that two types of visits, neurological and gastrointestinal, were predictive of student performance. Taken together, these results suggest that when students are suffering from ill health their academic performance is negatively impacted. Implications for improving health equity through data-driven educational interventions are discussed.

**Citation:** Rattermann MJ, Angelov A, Reddicks T, Monk J (2021) Advancing health equity by addressing social determinants of health: Using health data to improve educational outcomes. PLoS ONE 16(3): e0247909. <https://doi.org/10.1371/journal.pone.0247909>

**Editor:** Tamara Taggart, George Washington University, UNITED STATES

**Received:** March 5, 2020; **Accepted:** February 16, 2021; **Published:** March 17, 2021

**Copyright:** © 2021 Rattermann et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability:** The data that support the findings of this study are openly available. All data files are available from the Mendeley database in “Paramount Health Data Project 2020”, V1, (doi: [10.17632/tcbh2kzhh5.1](https://doi.org/10.17632/tcbh2kzhh5.1)).

**Funding:** This work was funded with grants from the Community Health Network Foundation ([www.ecommunity.com/community-foundation](http://www.ecommunity.com/community-foundation)) and Prosperity Indiana ([www.prosperityindiana.org](http://www.prosperityindiana.org)) to the Paramount Health Data Project. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Health equity means that everyone has a fair and just opportunity to be as healthy as possible [1]. To achieve health equity, access to the conditions and resources that positively influence health must be available to everyone, including historically excluded or marginalized groups of society. These conditions and resources that influence health are often referred to as the *social determinants of health*. Specifically, social determinants of health are conditions in the environment in which people live, learn, work, play, worship, and age that affect a wide range of health, functioning, and quality of life outcomes and risks. Examples of social determinants of health include quality of education, public safety, access to health care services, social support, residential segregation, among others [2].

These social determinants of health often interact with one another to create complex systems that influence long-term outcomes [3, 4]. Two such factors that have a significant impact on children of poverty in the United States are access to health care and quality of education. Equitable access to quality education in the United States is one of the areas that has the most striking disparities based on socio-economic status. Economic segregation, as measured by how evenly poor and nonpoor students are distributed among public schools, has been on the increase [5], with students of poverty being more likely to be in schools that lack access to the high-quality resources and learning opportunities that are often available to their wealthier peers [6, 7]. For example, teachers at high poverty schools have less experience than teachers in low poverty schools [8, 9] and often have fewer formal qualifications and have less resources to use during classroom instruction [10]. This inequity directly impacts student academic achievement, with students in high-poverty/high-minority schools consistently performing poorer in measures of academic success than white students in low-poverty schools [7, 11].

Access to quality health care is also affected by socioeconomic status. Despite the great gains made in access to healthcare since the passage of the Affordable Care Act, children from economically disadvantaged communities often lag behind their peers in more affluent communities in access to quality health care [12–14]. Over one quarter (26.3%) of poor children under 18 years of age in the United States have gone more than 6 months without having contact with a doctor or other health care professional, with 4.2% of those children having no contact with a doctor or health care professional in over five years [15].

Many schools are addressing the lack of access to healthcare through school nurse programs and school-based health centers. School-based health care provides access to primary and preventive health care services for students who are at risk for negative medical, social and academic outcomes [16, 17]. For children of poverty, the school nurse and the school-based health center are often the only medical care that is available to them [18, 19]. School based health care also has the added benefit of being located where children spend large periods of time—often 8 or more hours a day and over 180 days per year. Consequently, there is no need for parents to provide transportation to the location of the health provider, or for them to take time off work to accompany their child to the doctor [20–22] leading to more frequent access to health care [23, 24]. Additionally, school nurses and school-based health centers are often located in schools that serve students of poverty who are more likely to be chronically stressed, tired, hungry, and have vision and hearing problems, all factors that put a student at risk for academic failure [25–27]. School-based health care has been shown to have a positive effect on students, both in their academic behaviors such as attendance, suspensions, withdrawals from school and graduation rates [28–31] and in their social and emotional well-being, with a positive effect on school connectedness [32–35] and on quality of life [36].

The goal of this research is to examine the relationship between two social determinants of health: access to health care and quality education. The data for this research comes from a high-performing K-8 charter school that serves a high poverty community with a large minority population; 65% of the students in the school are African-American or Hispanic and 83% are children of poverty (as measured by free and reduced lunch). The primary hypothesis of this research is that academic achievement, as measured by standardized test scores, will be impacted by overall student health, as measured by the number of visits to a school nurse. A secondary hypothesis is that the reason for a visit to the school nurse, not just the number of visits, will also have an impact. Specifically, that respiratory complaints or complaints of parasites and skin rashes, both of which previous research has found to be related to poverty, [37, 38] will be found to impact performance on measures of academic achievement.

## Materials and methods

### Ethics statement

Prior to commencement of the study, ethical approval (IRB No.2) was obtained from the Institutional Review Board (IRB) of IntegReview IRB. The research was determined to be a clinical investigation not involving greater than minimal risk under an expedited review. Assent was waived under 45 CFR, 46.116(d) due to the use of retrospective data that were deidentified for this research.

### Participants

The data used in these analyses are from the Paramount School of Excellence Brookside (Paramount Brookside), which is a public charter school on the near eastside of Indianapolis. Paramount Brookside serves urban students in grades K-8. With a diverse student body of more than 1000 students, Paramount Brookside provides a sample of students who are living in poverty—with 83% of students qualifying for free and reduced meals, a common indicator of poverty. The students at Paramount Brookside are also ethnically diverse, with African American students making up 49% of Paramount Brookside's population, while Latinos are 16%, Multiracial students are 10%, and European American students are 25%. Almost 20% of PSOE students are identified as students with disabilities and receive special education services and 6% are identified as English Language Learners. Paramount Brookside

is a high performing school, with approximately 90% of all Paramount Brookside students passing the state mandated IREAD 3 test and more than 80% of all Paramount Brookside students passing both the English and Math portions of the state mandated ISTEP test during the years corresponding to the data being used in this research.

To examine the questions posed in this study a dataset was constructed that included student demographics such as gender, race/ethnicity and grade, the number of visits to the school nurse, and student scores on Acuity, a standardized assessment of student academic achievement. The Acuity assessment is administered longitudinally at three time points of the year (i.e., Acuity A is given at the beginning of the academic year, Acuity B is given in the middle, and Acuity C is given near the end) in two subject areas: Mathematics and English Language Arts (ELA).

#### Data collection method

Since 2013 Paramount Brookside has employed a full-time nurse, who during the time of the data collection saw 70–100 students per day. These visits range from a student who is sick with the flu to a student who find the class overwhelming and simply needs to talk to a sympathetic adult. The data were entered into the University of Tennessee's Consortium for Health Education, Economic Empowerment, and Research (CHEER) data collection platform, following the CHEER protocols for all visits to the nurse's office. The student health data and the academic achievement data were deidentified and merged by an independent third party acting as an honest broker of the data. All visits to the school nurse for either administration of prescription or over the counter drugs were removed from the dataset. All reported analyses were performed on the deidentified dataset.

#### Categorization of variables

At the time of a student visit to the school nurse, the student characteristics (e.g., grade, gender and race/ethnicity) were recorded using the CHEER data collection platform. The school nurse also recorded the reason for the student's visit and the outcome of that visit (e.g., the student was sent back to class, was sent home, etc.). The school nurse categorized the reason for the student visit into one of 16 categories (excluding regular prescription and over the counter medication administrations) from the CHEER protocol: (1) Cardiovascular (e.g., blood pressure check, chest pain), (2) Dental (e.g., toothache, lost tooth), (3) Dermatological (e.g., rash, eczema, bug bite, splinter), (4) Eye/Ears/Nose/Throat (e.g., earache, sore throat, pinkeye), (5) Endocrine (e.g., blood sugar check), (6) Gastrointestinal (e.g., heartburn, nausea, diarrhea), (7) Genitourinary (e.g., urinary frequency, soiled clothing), (8) Gynecological/Obstetrical (e.g., menstrual cramps), (9) Musculoskeletal (e.g., body aches, swelling), (10) Nutrition/Metabolic (G tube feed), (11) Neurological (e.g., headache, dizziness, falling asleep in class), (12) Parasites/Infections (e.g., bed bugs, head lice, ticks), (13) Disorder from Physical Agents (e.g., electrical shock), (14) Psychosocial (e.g., anxiety, altercation, tearful, needs to talk), (15) Respiratory (e.g., asthma, choking, cough, shortness of breath), (16) Other/Miscellaneous (e.g., physical injury, chills, illness).

The data from the 2013–14, 2015–16 and 2016–17 school years were included in these analyses. The data from the 2014–2015 school year were not included due to the lack of student achievement data for that year. 584 students from grade 3–8 contributed 933 sets of data (one year of test scores and one year of health data) to this dataset. Of the 933 sets of data, 48% (or 452) were contributed by girls and 52% (481) were contributed by boys. The majority of the datasets were contributed by African-American students (45% or 419 datasets), followed by white students (29% or 272 datasets), Hispanic students (17% or 157 datasets) and then multiracial students (9% or 86 datasets). Each of these students provided at least one year of Acuity data. Overall, these

students visited the nurse's office a total of 4467 times over the 3 years of the data being used in this study. The most common frequency of nurse office visits per student was "none" for 21% of students, followed by 1–4 for 42% of students, and 5–10 for 30% of students. The remaining 7% of students visited the school nurse office from 11–84 times.

## Results

The quantitative data were analyzed using SPSS 26.0 (IBM Corp., Armonk, NY, USA). Participants' general characteristics were analyzed with descriptive statistics, significant differences between these general characteristics were analyzed with ANOVA. Post hoc analyses were done using the Bonferroni adjustment to correct for multiple comparisons. For correlations between the number of visits to the school nurse and educational outcomes, Pearson's correlation coefficients were computed.

### Results by demographics

Across the three school years, male students had slightly more yearly visits to the nurse (51%) as compared to female students (49%), with the mean number of visits to the school nurse per year for female students (5.31) and male students (5.17) roughly equal. African-American students accounted for the majority of the total nurse visits (50%) as compared to 32% for white students, 10% for Hispanic students and 8% for multiracial students. The mean number of visits to the school nurse was 5.83 for African-American students, 5.72 for White students, 3.29 for Hispanic students and 4.31 for Multiracial students.

Visit frequencies of students across grades varied significantly, with younger students accounting for more visits to the school nurse than older students. Specifically, students in grades 3 and 4 both accounted for 21% of visits to the school nurse, 5th grade students accounted for 18%, 6th grade students accounted for 16%, 7th grade students accounted for 14% and 8th grade students accounted for 9%. The mean number of visits by grade revealed that 3rd grade students visited the school nurse an average of 5.11 times, 4th grade students visited the school nurse an average of 6.38 times, 5th grade students visited an average of 5.55 times, 6th grade students visited an average of 5.39 times, 7th grade students visited an average of 3.52 times, and finally 8th grade students visited an average of 4.59 times.

These findings were confirmed by a race (Black, White, Hispanic and Multiracial) Gender (male and female) x Grade (3rd, 4th, 5th, 6th, 7th and 8th grades) within subjects Analysis of Variance performed on the total number of visits to the school nurse which found that Race was significant ( $F(3,884) = 4.54, p < .01$ ) as was Grade ( $F(3,885) = 4.54, p < .004$ ). There were no other main effects and no significant interactions. Bonferroni post hoc analyses revealed that the mean number of visits to the school nurse by African American (5.83) and White students (5.72) were not significantly different from each other but were significantly different from the mean number of visits to the school nurse by Hispanic (3.29) and Multiracial students (4.3) who were not significantly different from each other. Post hocs revealed a significant difference between the mean number of visits to the school nurse between 4th grade students (6.38) and 7th grade students (3.52).

### Correlation between health data and academic data

The primary hypothesis tested in this research is that the number of visits to the school nurse is predictive of student achievement scores. To determine the effect of the number and reason for visits to the school nurse office a series of bivariate correlations were performed over the 16 categories of health reasons and the three yearly Acuity scores for Language Arts and Mathematics. A conservative  $\alpha$  value of 0.01 was used to address the possibility of finding a significant result by chance alone because of multiple comparisons.

Pearson correlation analyses investigating the impact of the type and number of visits to the school nurse on academic achievement revealed a negative correlation between total number of visits and scores on the Acuity tests across time in Language Arts, with the comparison between the first administration in a school year of Language Arts Acuity A revealing a significant correlation (-.196) between the total number of visits to the school nurse and the Acuity scores. This relationship was also found in the second administration in a school year of Language Arts Acuity B, with the comparison between the second administration and the total number of visits to the school nurse also revealing a significant correlation (-.174). A weaker, but still significant correlation was also found between the third administration of Language Arts Acuity C (-.118). Similar results were found in Acuity Mathematics, although the correlations were not as strong as those with Language Arts. All of the significant correlations can be found in [Table 1](#) below.

**Table 1. Correlations between health data and academic achievement.**

<https://doi.org/10.1371/journal.pone.0247909.t001>

In addition to significant correlations between the total number of visits to the school nurse, significant correlations were found between Language Arts Acuity and Neurological, Gastrointestinal, Dermatological, Musculoskeletal, Eyes, Ears, Nose & Throat and “other” types of visits to the school nurse. These correlations were particularly strong for the first and second administration of Language Arts Acuity.

To further explore the relationship between the number and type of visits to the school nurse and academic achievement as measured by Acuity a series of fixed effects regressions, or linear regressions with unit and time fixed effects, were performed on the longitudinal academic assessment data. The use of a fixed effects regression also adjusted for unobserved unit-specific and time-specific confounds that may have been present in the data.

Because the data from the 2014–15 school year were not available, only the data from the 2015–16 and 2016–17 school years were contiguous. Consequently, these analyses were performed over a subset of the data that included only student data from those years. This decreased the dataset from 584 to 155 students. In this new dataset students from grade 3–8 contributed 155 sets of data (two year of test scores and two years of health data each). Of the 155 sets of data, 45% (or 141) were contributed by

girls and 55% (169) were contributed by boys. The majority of the datasets were contributed by African-American students (48% or 149 datasets), followed by white students (25% or 79 datasets), Hispanic students (17% or 54 datasets) and multiracial students (9% or 28 datasets).

The fixed effects regressions were performed with *student* as the unit variable and *year (1 and 2)* as the time variable. The outcome variable for these analyses was student *Growth Scores* for the Acuity English Language Arts and Mathematics assessments. Growth Scores are computed by subtracting student scores on the first Acuity administration in each academic year (Acuity A) from the final administration in each academic year (Acuity C) in each area (English Language Art and Mathematics). As a Student Growth Assessment, Acuity is designed to be used to determine student growth over the course of the academic year. Because the independent variable in these analyses (number/type of visits to the school nurse) is cumulative for the school year, student academic growth measured at the end of the academic year was determined to be the appropriate outcome variable.

The quantitative data were analyzed using Stata/IC 16.1 for Mac (StataCorp, College Station, TX, USA). All of the fixed effects (within) regressions were performed with *student* as the Panel variable and *Year (1 and 2)* as the Time variable. Hausman tests for random effects versus fixed effects models were performed for each of the significant analyses reported and in each case the null hypothesis was rejected, confirming that a fixed effects model was appropriate.

The Language Arts Acuity Growth analyses revealed significant effects of Total Number of Visits ( $F(1,154) = 6.38, p < .01$ ), Dermatological visits ( $F(1,154) = 4.53, p < .05$ ), and Gastrointestinal visits, ( $F(1,154) = 3.73, p < .05$ ). There were no significant effects revealed in the Mathematics Acuity Growth analyses.

#### Types of visits to the school nurse

The most common type of visits to the school nurse were for Gastrointestinal complaints (e.g., upset stomach, gastric reflux) and Ear, Nose and Throat (e.g., sore throats) which both accounted for 20% of the visits to the school nurse. Dermatological (e.g., rashes) at 16% of the total and Neurological (headaches) at 13% of the total were the two next most common complaints. The frequency and type of visits to the school nurse can be found in [Table 2](#) below.

**Table 2. Frequency of types of visits to the school nurse.**  
<https://doi.org/10.1371/journal.pone.0247909.t002>

## Discussion

The results of this research revealed that there is a significant negative correlation between the number of visits to a school nurse and academic achievement as measured by a longitudinal standardized assessment of academic achievement. This effect was stronger for English Language Arts than for Mathematics, and for English Language Arts this effect was consistent over three testing periods. Further analyses revealed that of the 16 different types of visits to the school nurse, performance in English Language Arts was correlated to neurological, gastrointestinal, dermatological, musculoskeletal, Eyes, Ears, Nose & Throat, and “other” categories of visits types.

Additional fixed effect regression analyses confirmed and expanded the results of the correlations, suggesting that there is a causal relationship between the number and type of visits to the school nurse and academic achievement. The fixed effects regression analyses found significant effects of the total number of visits to the school nurse, as well as the type of visits (dermatological and neurological) to the school nurse on student growth in Acuity scores in English Language Arts.

Overall, the results of this research support our primary hypothesis that academic achievement, as measured by standardized test scores, is impacted by overall student health, as measured by the number of visits to a school nurse. The results also support our secondary hypothesis that the reason for a visit to the school nurse, not just the number of visits, will also have an impact, although the specific category of complaints predicted based on the literature, parasites and respiratory complaints were not the categories found to be significant in this research. Rather, dermatological and gastrointestinal complaints were predictive of academic performance for the sample of students studied here. This particular finding may be due to characteristics of this population and deserves further study. It is also important to note that the categories used by the CHEER protocol are broad and the specific diagnoses (e.g, rash versus bug bite for dermatological and heartburn versus nausea for gastrointestinal) were not available for research and could have yielded more specific conclusions.

There are several possible explanations for the results of this research. First, the data suggests that students who are frequent visitors to the school nurse are simply unhealthy and frequently do not feel well during the school day, and this is impacting their ability to learn. As noted by Basch and his colleagues “healthier students are better learners” [39–41]. Research on the impact of health on academic behaviors supports this argument [39, 40, 42–44]. Children who are unhealthy are at higher risk for dropping out [45], failing classes [46], and grade retention [44], while children who exercise regularly, are at a healthy body weight and have access to healthy diets show increased cognitive abilities and positive academic behaviors [42].

A second possible explanation is that students who frequent the school nurse are missing valuable class time and are consequently performing poorly on the standardized assessments. This explanation may be valid when considering the students who have visited the school nurse multiple time per week, but the majority of students in the sample visited the school nurse under 10 times in an academic year. With the average time spent with a school nurse in this sample being approximately 5 minutes, even with travel time to and from the classroom the amount of instruction time missed is minimal in the context of the larger school year.

What is more likely is that many students who are frequent visitors to the school nurse are exhibiting symptoms of larger stressors in their lives. The data from this research supports this hypothesis, with stress related gastrointestinal and neurological complaints making up nearly one third of the reasons for visits to the school nurse, and gastrointestinal complaints also being one of the two significant categories found by the regression analyses. The most common types of somatic complaints (where students visit the school nurse because of vague, varied and sometimes unsubstantiated complaints) found in the literature are headaches, followed



closely by stomachaches [47–51]. These somatic complaints have been found to be related to comorbid mental health problems with children classified as “somatizers” being four times more likely to screen for mental health problems than typical peers [52]. Further, children who exhibit somatic complaints are more often in minority, urban and non-intact and less educated families [52].

This research has demonstrated that there is a significant relationship between student health and academic achievement. Additional research needs to be done to provide more specific information regarding the interactions between student health and performance in school. The current findings, however, are enough to begin the process of using data from one social determinant of health—access to health care—to address another—access to a quality education. Based on the significant negative correlation between visits to the school nurse and scores on academic assessments, Paramount School of Excellence Brookside has begun the process of integrating student health data into Multi-Tiered System of Supports (MTSS) [53]. MTSS is a systemic, continuous-improvement framework in which student data is used to inform decision-making that supports students in their academic and socio-emotional development. Specifically, in the MTSS process student data from standardized assessments are used to determine whether a student’s needs are being met by the Tier 1 interventions that are implemented with all students in the classroom setting or if Tier 2 services, such as small group tutoring, are needed to ensure student success. If the student does not show progress with the Tier 2 interventions, then Tier 3 supports will be provided, often through Special Education or other services. MTSS is most effective when it is an early intervention—the sooner a student receives the MTSS services the more effective they will be. Paramount Brookside is using student health data to trigger the MTSS process much sooner than would normally be the case if the move to Tier 2 was triggered by the results of standardized academic assessments given in the middle of the year.

The ultimate goal of health equity is that everyone has a fair and just opportunity to be as healthy as possible. Addressing social determinants of health in childhood can lead to positive changes across the lifespan, with access to education impacting yearly mortality rates and length of life in adults [54, 55]. In fact, high school graduates live an average of six years longer than adults with less than a high school education, regardless of gender or racial/ethnicity group [56, 57], and have a higher quality of life, reporting greater well-being, fewer chronic health problems, and fewer common illnesses such as colds, headaches and body aches [58]. The results of this research point the way to better use of student health data in schools, which will lead to both better educational and better health outcomes for everyone.

## Acknowledgments

The Authors would like to thank the staff and students of Paramount School of Excellence Brookside, The M.A. Rooney Foundation, the Collaborative Research Institute at Indiana University, the Community Hospital Foundation and Prosperity Indiana.

## References

1. Braveman P, Arkin E, Orleans T, Proctor D, Plough A, Austin D, et al. What Is Health Equity? And What Difference Does a Definition Make? National Collaborative for Health Equity MBA, Evidence for Action. Robert Wood Johnson Found [Internet]. 2017;(May):20. Available from: [http://www.rwjf.org/content/dam/farm/reports/issue\\_briefs/2017/rwjf437393](http://www.rwjf.org/content/dam/farm/reports/issue_briefs/2017/rwjf437393)  
[View Article](#) • [Google Scholar](#)

2. Healthy People 2020 [Internet]. Washington, DC US; [cited 2020 Feb 9]. Available from: <https://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-of-health>
3. Ahnquist J, Wamala SP, Lindstrom M. Social determinants of health—a question of social or economic capital? Interaction effects of socioeconomic factors on health outcomes. *Soc Sci Med*. 2012 Mar;74(6):930–9. Epub 2012 Jan 21. pmid:22305807.  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
4. Jayasinghe S. Social determinants of health inequalities: towards a theoretical perspective using systems science. *Int J Equity Health* 14, 71 (2015). pmid:26303914  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
5. Owens A. Income Segregation Between School Districts and Inequality in Students' Achievement. *Soc of Educ*. 2018;91(1):1–27.  
[View Article](#) • [Google Scholar](#)
6. Owens A, Reardon SF, Jencks C. Income Segregation Between Schools and School Districts. *Am Educ Res J*. 2016;53(4):1159–97.  
[View Article](#) • [Google Scholar](#)
7. Lacour M, Tissington LD. The effects of poverty on academic achievement. *Educ Res Rev*. 2011;6(7):522–7.  
[View Article](#) • [Google Scholar](#)
8. Clotfelter C, Ladd H, Vigdor J, Wheeler J. High-poverty schools and the distribution of teachers and principals. *North Carol Law Rev*. 2007;85(5):1345–79.  
[View Article](#) • [Google Scholar](#)
9. Hanushek EA, Rivkin SG. The Distribution of Teacher Quality and Implications for Policy. *Annu Rev Econom*. 2012;4(1):131–57.  
[View Article](#) • [Google Scholar](#)
10. Phillips M, Chin T. School inequality: What do we know? *Soc Inequal*. 2004;467–519.  
[View Article](#) • [Google Scholar](#)
11. Musu-Gillette L, de Brey C, McFarland J, Hussar W, Sonnenberg W, Wilkinson-Flicker S. Status and trends in the education of racial and ethnic groups 2017 (NCES 2017–051). Vol. 1. Washington, DC; 2017.
12. Larson K, Cull WL, Racine AD, Olson LM. Trends in access to health care services for US children: 2000–2014. *Pediatrics*. 2016;138(6). pmid:27940710  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

13. Chen J, Vargas-Bustamante A, Mortensen K, Ortega AN. Racial and ethnic disparities in health care access and utilization under the affordable care act. *Med Care*. 2016;54(2):140–6. pmid:26595227  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
14. Flores G. Racial and ethnic disparities in the health and health care of children. *Pediatrics*. 2010;125(4). pmid:20351000  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
15. National Center for Health Statistics. National Health Interview Survey. 2018.
16. Gustafson EM. History and overview of school-based health centers in the US. *Nurs Clin North Am*. 2005;40(4):595–606. pmid:16324934  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
17. Strolin-Goltzman J. The relationship between school-based health centers and the learning environment. *J Sch Health [Internet]*. 2010 Mar;80(3):153–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20236418> pmid:20236418  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
18. Albright K, Barnard J, O’Leary S, Federico S, Saville A, Lockhart S, et al. School-Based Health Centers as Medical Homes: Parents’ and Adolescents’ Perspectives. *Acad Pediatr*. 2016 May 1;16(4):381–6. pmid:26329018  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
19. O’Leary ST, Lee M, Federico S, Barnard J, Lockhart S, Albright K, et al. School-based health centers as patient-centered medical homes. *Pediatrics*. 2014;134(5):957–64. pmid:25287455  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
20. Anglin TM, Naylor KE, Kaplan DW. Comprehensive school-based health care: High school students’ use of medical, mental health, and substance abuse services. *Pediatrics [Internet]*. 1996 Mar [cited 2011 Oct 1];97(3):318–30. pmid:8604264  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
21. Barnett SE, Allison M, Roland MM, Devore CD, Ancona R, Gunther R, et al. School-based health centers and pediatric practice. Vol. 129, *Pediatrics*. 2012. p. 387–93. pmid:22291117  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
22. Brindis CD, Klein J, Schlitt J, Santelli J, Juszczak L, Nystrom RJ. School-based health centers: Accessibility and accountability. *J Adolesc Heal*. 2003;32(6 SUPPL.):98–107. pmid:12782448  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

23. Arenson M, Hudson PJ, Lee N, Lai B. The Evidence on School-Based Health Centers: A Review. *Glob Pediatr Heal*. 2019;6:2333794X1982874. pmid:30815514  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
24. Brindis CD. The “state of the state” of school-based health centers: Achieving health and educational outcomes. *Am J Prev Med* [Internet]. 2016;51(1):139–40. Available from: pmid:27320218  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
25. Adelman HS, Barker LA, Nelson P. A Study of a School-Based Clinic: Who Uses It and Who Doesn't? *J Clin Child Psychol*. 1993;22(1):52–9.  
[View Article](#) • [Google Scholar](#)
  
26. Knopf JA, Finnie RKC, Peng Y, Hahn RA, Truman BI, Vernon-Smiley M, et al. School-based health centers to advance health equity: A community guide systematic review. *Am J Prev Med*. 2016 Jul 1;51(1):114–26. pmid:27320215  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
27. Kisker EE, Brown RS. Do school-based health centers improve adolescents' access to health care, health status, and risk-taking behavior? *J Adolesc Heal* [Internet]. 1996 May [cited 2011 Oct 2];18(5):335–43. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9156546> pmid:9156546  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
28. Gall G, Pagano M, Desmond M, Perrin J, Murphy J. Utility of Psychosocial Screening at a SBHC. *J Sch Health*. 2000;70(7):292–8. pmid:10981284  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
29. Kerns SEU, Pullmann MD, Walker SC, Lyon AR, Cosgrove TJ, Bruns EJ. Adolescent use of school-based health centers and high school dropout. *Arch Pediatr Adolesc Med*. 2011 Jul;165(7):617–23. pmid:21383256  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
30. McCord MT, Klein JD, Foy JM, Fothergill K. School-based clinic use and school performance. *J Adolesc Heal*. 1993;14(2):91–8. pmid:8476879  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
31. Walker SC, Kerns SEU, Lyon AR, Bruns EJ, Cosgrove TJ. Impact of School-Based Health Center Use on Academic Outcomes. *J Adolesc Heal* [Internet]. 2010 Mar [cited 2011 Sep 26];46(3):251–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20159502>  
[View Article](#) • [Google Scholar](#)
  
32. Bersamin M, Coulter RWS, Gaarde J, Garbers S, Mair C, Santelli J. School-Based Health Centers and School Connectedness. *J Sch Health*. 2019 Jan 1;89(1):11–9. pmid:30506695

[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

33. Strolin-Goltzman J, Sisselman A, Melekis K, Auerbach C. Understanding the relationship between school-based health center use, school connection, and academic performance. *Heal Soc Work*. 2014;39(2):83–91.  
[View Article](#) • [Google Scholar](#)
34. Whitaker K. [Internet]. ProQuest Dissertations and Theses. 2015 [cited 2019 Nov 26].
35. Centers for Disease Control and Prevention. School connectedness: Strategies for increasing protective factors among youth. US Dep Heal Hum Serv [Internet]. 2009 [cited 2019 Dec 8];1–24. Available from: [www.cdc.gov/HealthyYouth](http://www.cdc.gov/HealthyYouth).  
[View Article](#) • [Google Scholar](#)
36. Wade TJ, Mansour ME, Line K, Huentelman T, Keller KN. Improvements in Health-Related Quality of Life Among School-Based Health Center Users in Elementary and Middle School. *Ambul Pediatr*. 2008;8(4):241–9. pmid:18644546  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
37. Hotez PJ. Neglected diseases and poverty in “The other America”: The greatest health disparity in the United States? *PLoS Negl Trop Dis*. 2007;1(3):1–3. pmid:18160982  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
38. Francis L, DePriest K, Wilson M, Gross D. Child poverty, toxic stress, and social determinants of health: Screening and care coordination. *Online J Issues Nurs* [Internet]. 2018;23(3). Available from: pmid:31427855  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
39. Basch CE. Healthier Students Are Better Learners: A Missing Link in School Reforms to Close the Achievement Gap. *Equity Matters. Research Review No. 6*. Campaign Educ Equity, Teach Coll Columbia Univ [Internet]. 2010;(6):1–109.  
[View Article](#) • [Google Scholar](#)
40. Basch CE. Healthier Students Are Better Learners: High-Quality, Strategically Planned, and Effectively Coordinated School Health Programs Must Be a Fundamental Mission of Schools to Help Close the Achievement Gap. *J Sch Health*. 2011;81(10):650–62. pmid:21923878  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
41. Michael SL, Merlo CL, Basch CE, Wentzel KR, Wechsler H. Critical Connections: Health and Academics. *J Sch Health*. 2015;85(11):740–58. pmid:26440816  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)

42. Shaw SR, Gomes P, Polotskaia A, Jankowska AM. The relationship between student health and academic performance: Implications for school psychologists. *Sch Psychol Int* [Internet]. 2015 Apr 30 [cited 2019 Nov 26];36(2):115–34. Available from: <http://journals.sagepub.com/doi/10.1177/0143034314565425>  
[View Article](#) • [Google Scholar](#)
  
43. Scheier LM, Botvin GJ, Griffin KW, Diaz T. Latent growth models of drug refusal skills and adolescent alcohol use. *J Alcohol Drug Educ* [Internet]. 1999 Mar;44(3):21–48.  
[View Article](#) • [Google Scholar](#)
  
44. Bradley BJ, Greene AC. Do health and education agencies in the united states share responsibility for academic achievement and health? A review of 25 years of evidence about the relationship of adolescents' academic achievement and health behaviors. *J Adolesc Heal* [Internet]. 2013;52(5):523–32. Available from: <http://dx.doi.org/10.1016/j.jadohealth.2013.01.008>  
[View Article](#) • [Google Scholar](#)
  
45. Vaughn MG, Salas-Wright CP, Maynard BR. Dropping out of school and chronic disease in the United States. *J Public Health (Bangkok)* [Internet]. 2014;22(3):265–70. Available from: [pmid:25232516](https://pubmed.ncbi.nlm.nih.gov/25232516/)  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
46. Needham BL, Crosnoe R, Muller C. Academic Failure in Secondary School: The Inter-Related Role of Health Problems and Educational Context. *Soc Probl.* 2004;51(4):569–86. [pmid:20354573](https://pubmed.ncbi.nlm.nih.gov/20354573/)  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
47. Bernstein G. A., Massie E. D., Thuras P. D., Perwien A. R., Borchardt C. M., & Crosby RD. Somatic symptoms in anxious-depressed school refusers. *J Am Acad Child Adolesc Psychiatry.* 1997;36:661–8. [pmid:9136501](https://pubmed.ncbi.nlm.nih.gov/9136501/)  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
48. Egger HL, Costello EJ, Erlanli A, Angold A. Somatic Complaints and Psychopathology in Children and Adolescents: Stomach Aches, Musculoskeletal Pains, and Headaches. *J Am Acad Child Adolesc Psychiatry* [Internet]. 1999 Jul 1;38(7):852–60. Available from: [pmid:10405503](https://pubmed.ncbi.nlm.nih.gov/10405503/)  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
49. Lieb R, Zimmermann P, Friis RH, Hofler M, Tholen S, Wittchen HU. The natural course of DSM-IV somatoform disorders and syndromes among adolescents and young adults. a Prospect Community Study [Internet]. 2013;1–14. Available from: [http://www.qucosa.de/fileadmin/data/qucosa/documents/11001/578\\_PP.pdf%5Cnpapers3://publication/uuid/82EF5F2A-CFCD-4E0A-9B32-AF71A332CCF9](http://www.qucosa.de/fileadmin/data/qucosa/documents/11001/578_PP.pdf%5Cnpapers3://publication/uuid/82EF5F2A-CFCD-4E0A-9B32-AF71A332CCF9)  
[View Article](#) • [Google Scholar](#)

50. Natvig GK, Albrektsen G, Anderssen N, Qvarnström U. School-related Stress and Psychosomatic Symptoms Among School Adolescents. *J Sch Health* [Internet]. 1999;69(9):362–8. Available from: pmid:10633321  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
51. Schneider MB, Friedman SB, Fisher M. Stated and unstated reasons for visiting a high school nurse's office. Vol. 16, *Journal of Adolescent Health*. Netherlands: Elsevier Science; 1995. p. 35–40.
  
52. Campo JV, Jansen-McWilliams L, Comer DM, Kelleher KJ. Somatization in Pediatric Primary Care: Association With Psychopathology, Functional Impairment, and Use of Services. *J Am Acad Child Adolesc Psychiatry* [Internet]. 1999 Sep 1;38(9):1093–101. Available from: pmid:10504807  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
53. Angelov AD, Pettinga D, Bateman D. The Paramount Health Data Project. In: *Hashtags & Headlines Marketing for School Leaders*. New York, NY US: Roman & Littlefield; 2020. p. 82–5.
  
54. Feinglass J, Lin S, Thompson J, Sudano J, Dunlop D, Song J, et al. Baseline health, socioeconomic status, and 10-year mortality among older middle-aged Americans: Findings from the health and retirement study, 1992–2002. *Journals Gerontol—Ser B Psychol Sci Soc Sci*. 2007;62(4):209–17. pmid:17673534  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
55. Steenland K, Henley J, Calle E, Thun M. Individual- and area-level socioeconomic status variables as predictors of mortality in a cohort of 179,383 persons. *Am J Epidemiol*. 2004;159(11):1047–56. pmid:15155289  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
56. Hummer RA, Hernandez EM. Population bulletin: The effect of educational attainment on adult mortality in the United States. *Popul Bull*. 2013;68(1):1–20. pmid:25995521  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
57. Krueger PM, Tran MK, Hummer RA, Chang VW. Mortality attributable to low levels of education in the United States. *PLoS One*. 2015;10(7):1–13. pmid:26153885  
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
  
58. Ionescu DD, Ionescu AM, Jaba E. The investments in education and quality of life. *J Knowl Manag Econ Inf Technol Spec*. 2013;(December):141–58.  
[View Article](#) • [Google Scholar](#)